

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

BC

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 :

A61K 43/00, 49/02
C07D 451/00, 451/02

A1

(11) International Publication Number:

WO 93/09814

(43) International Publication Date:

27 May 1993 (27.05.93)

(21) International Application Number: PCT/US92/09482

(22) International Filing Date: 13 November 1992 (13.11.92)

(30) Priority data:
792,648

15 November 1991 (15.11.91) US

(71) Applicants: RESEARCH TRIANGLE INSTITUTE [US/US]; Office of Research Contracts, 3040 Cornwallis Road, Research Triangle Park, NC 27709 (US). NATIONAL INSTITUTES OF HEALTH [US/US]; 9000 Rockville Pike, Bethesda, MD 20892 (US).

(72) Inventors: KUHAR, Michael, J. ; 2415 Briarwood Road, Baltimore, MD 21209 (US). BOJA, John, W. ; 1 Castell Court, Baltimore, MD 21236 (US). CARROL, Frank, Ivy ; 5124 Willowhaven Drive, Durham, NC 27712 (US). LEWIN, Anita, H. ; 804 Cedar Falls Road, Chapel Hill, NC 27515 (US). ABRAHAM, Philip ; 310 Homestead Drive, Cary, NC 27511 (US).



(74) Agents: KELBER, Steven, B. et al.; Oblon, Spivak, McClelland, Maier & Neustadt, Fourth Floor, 1755 South Jefferson Davis Highway, Arlington, VA 22202 (US).

(81) Designated States: AU, CA, FI, JP, KR, NO, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE).

Published

With international search report.

(54) Title: COCAINE RECEPTOR BINDING LIGANDS



(57) Abstract

Novel compounds show high affinity for specific cocaine receptors in the brain, particularly dopamine transporter sites, and have formula (I), wherein Y = CH₂R₃, CO₂R₂ or (II), R₁ = hydrogen, C₁₋₅ alkyl, R₂ = hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₄ alkoxy, C₁₋₆ alkynyl, halogen or amine, R₃ = OH, hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₄ alkoxy, Cl, Br, I, CN, NH₂, NHC₁₋₆ alkyl, NC₁₋₆ alkyl, OCOC₁₋₆ alkyl, OCOC₁₋₃ alkylaryl, A = S, O or NH, X = H, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₄ alkoxy, C₁₋₆ alkynyl, halogen, amino, acylamido, and Z = H, I, Br, Cl, F, CN, CF₃, NO₂, N₃, OR₁, CO₂NH₂, CO₂R₁, C₁₋₆ alkyl, NR₄R₅, NHCOR₅, NHCO₂R₆, wherein R₄-R₆ are each C₁₋₆ alkyl.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

Description

Cocaine Receptor Binding Ligands

This application is a continuation-in-part application of U.S. Patent Application 07/564,755, filed August 9, 1990, and U.S. PCT Application PCT/US91/05553, filed August 9, 1991 (attorney docket 2025-055-27 PCT), filed in the U.S. PCT Receiving Office and designating the United States.

Technical Field

This invention is directed to a class of binding ligands for cocaine and other receptors in the brain. Specifically, a novel family of compounds shows high binding specificity and activity, and, in a radiolabeled form, can be used to bind to these receptors, for biochemical assays and imaging techniques.

Background Art

This application claims priority from U.S. Patent Application 07/564,755 and U.S. PCT Application PCT/US 91/05553, filed August 9, 1991, both applications being incorporated herein by reference. In U.S. Application Serial No. 07/564,755, there is disclosure of a family of compounds exhibiting particularly high specificity and affinity for cocaine receptors and other neurotransmitter receptors in the brain of the formula:


Where the broken line represents an optional chemical bond and the substituents at 2 and 3 may be at any position;

The iodo substituent may be at o, m, p, or multisubstituted;

$R_1 = \text{CH}_3, \text{CH}_2\text{CH}=\text{CH}_2, (\text{CH}_2)_n\text{C}_6\text{H}_5 \quad n = 1-4;$

-2-

$R_2 = \text{CH}_3, \text{C}_2\text{H}_5, \text{CH}_3(\text{CH}_2)_3, (\text{CH}_3)_2\text{CH}, \text{C}_6\text{H}_5, \text{C}_6\text{H}_5\text{CH}_2, \text{C}_6\text{H}_5(\text{CH}_2)_2;$
 $X = \text{pharmacologically acceptable anion}$



Sites of specific interest included cocaine receptors associated with dopamine transporter sites.

Subsequently, in the U.S. PCT Application from which priority is claimed, and which is incorporated herein by reference, the values for R_1 and R_2 were expanded, such that R_1 may be an alkyl of 1-7 carbon atoms, $\text{CH}_2\text{CR}_3=\text{CR}_4\text{R}_5$ wherein R_3 - R_5 are each, independently C_{1-6} alkyl, or phenyl compounds of the formula $\text{C}_6\text{H}_5(\text{CH}_2)_y$, wherein $y = 1-6$. The PCT filing also reveals the affinity of these compounds for cocaine receptors associated with serotonin transporters, and confirms, for the first time, that the in vitro binding reported in the earlier-filed application, is confirmed in in vivo testing. Specific disclosure for a variety of applications, including using the receptors in both PET and SPECT scanning, wherein either the iodine substituent, or one of the carbon groups is radioactive (I-123 , 125 or 131 and C11) thus providing methods for scanning for the presence of specific cocaine receptors appears. Such scanning processes may be used to determine physiological conditions, such as Parkinson's Disease, to examine in general the density and distribution of specific cocaine receptors in various parts of the brain and/or body, to determine the efficacy of neurological treatments aimed at halting or reversing the degeneration of specific nerves in the brain, and screening drugs, such as antidepressant drugs.


-3-

The affinity of these compounds, as reported in the applications incorporated, is surprisingly high, and compared with prior art compounds, such as [^3H]WIN 35,428, the novel compounds of these applications exhibit extremely low IC_{50} values for binding inhibition.


Other compounds exhibiting this type of binding activity and specificity would further advance this art, making it possible to obtain better scanning, with higher reliability, and lower levels.

Disclosure of the Invention

Applicants' invention resides in the discovery of a new family of compounds of the formula



Wherein $\text{Y} = \text{CH}_2\text{R}_3$, CO_2R_2 or



-4-

R_1 = hydrogen, C_{1-5} alkyl,

R_2 = hydrogen, C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-4} alkoxy, C_{1-6} alkynyl, halogen or amine,

R_3 = OH, hydrogen, C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-4} alkoxy, Cl, Br, I, CN, NH_2 , NHC_{1-6} alkyl, NC_{1-6} alkyl, $OCOC_{1-6}$ alkyl, $OCOC_{1-3}$ alkylaryl,

A = S, O or NH

X = H, C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-4} alkoxy, C_{1-6} alkynyl, halogen, amino, acylamido,

and

Z = H, I, Br, Cl, F, CN, CF_3 , NO_2 , N_3 , OR_1 , CO_2NH_2 , CO_2R_1 , C_{1-6} alkyl, NR_4R_5 , $NHCOR_5$, $NHCO_2R_6$,

wherein R_4 - R_6 are each C_{1-6} alkyl.

These compounds exhibit usually high affinity for binding to receptors for the dopamine transporter site, as well as the serotonin transporter site, again based on inhibition of [3H]WIN 35,428 binding. It will be immediately apparent that certain of the compounds embraced within this novel class exhibit iodine substituents, and thus can be made easily radioactive, by substitution of a radioactive iodine, according to the synthesis scheme set forth in the prior applications incorporated herein by reference. In those circumstances where no radioactive iodine is provided, the compounds may be made radioactive by selective use of at least one carbon that is radioactive, e.g., [^{11}C].

Best Mode for Carrying Out the Invention

The compounds of this invention can be prepared according to the synthesis methods described in the parent applications.

Additionally, specific synthesis routes and examples are set forth herein.

3 β -[3'-Iodo-4'-aminophenyl]tropan-2 β -carboxylic Acid Methyl Ester Dihydrochloride (1a). To a solution of 3 β -[4'-aminophenyl]tropan-2 β -carboxylic acid methyl ester (300 mg, 1.094 mmol) in glacial AcOH (15 mL) was added dropwise ICI (195 mg, 1.201 mmol) at room temperature for 3 h under N₂. After removal of solvent, the residue was diluted with H₂O, and then the pH was adjusted to basic with concentrated ammonia solution. The mixture was extracted with CHCl₃ which was washed with water and brine. After drying over MgSO₄, the solvent was evaporated to an oily product which was purified by flash chromatography (hexane-ether, 4:1). The collected fraction was converted to HCl salt with HCl/ether to yield 140 mg (29%) of 3 β -[3'-iodo-4'-aminophenyl]tropan-2 β -carboxylic acid methyl ester dihydrochloride (1a): mp 170-173°C; [α]_D²⁵ 90.9° (c 0.055, MeOH), ¹H NMR (250 MHz, CDCl₃) δ 1.65 (m, 3), 2.09 (m, 2), 2.2 (s, 3, NCH₃), 2.45 (m, 1), 2.75 (m, 1, H-2), 2.8 (m, 1, H-3), 3.33 (m, 1, H-5), 3.45 (m, 4, H-1, OCH₃), 3.95 (m, 2, NH₂), 6.65 (d, 1, J = 8.7, ArH), 7.05 (dd, 1, j = 8.7, J = 1.5, ArH), 7.42 (d, J = 1.5, 1, ArH).

Anal. Calcd for C₁₆H₂₁IN₂O₂·2 HCL·H₂O: C, 39.12; H, 5.13; N, 5.70. Found: C, 39.12, H, 5.16; N, 5.63.

3 β -[3'-Iodo-4'-azidophenyl]tropan-2 β -carboxylic Acid Methyl Ester Hydrochloride (1b). To a solution of 3, β -[3'-iodo-4'-aminophenyl]tropan-2 β -carboxylic acid methyl ester dihydrochloride (1a) (90 mg, 0.1902 mmol) in 1 mL of AcOH (3M) was added an aqueous solution of NaNO₂ (17.3 mg, 0.2661 mmol, in 0.5 mL of H₂O) at 0°C. After 30 min at this temperature NaN₃ (19 mg, 0.2754 mmol) in 0.5 mL of H₂O was added dropwise to the reaction mixture and stirred for 30 min at 0°C, then 30

-6-

min at room temperature. After removal of all solvent by evaporation, the residue was dissolved in CHCl_3 and washed with H_2O . The organic layer was dried over MgSO_4 and concentrated to give an oil which was converted to HCl salt to yield 64 mg (72.7%) of 3β -[3'-iodo-4'-azidophenyl]tropan- 2β -carboxylic acid methyl ester hydrochloride (1b) as a yellowish solid: mp 140 - 143°C ; $[\alpha]^{25}_{\text{D}}$ -97.4° (c 0.115), MeOH); ^1H NMR (250 MHz, CDCl_3) δ 1.51-1.73 (m, 3) 2.07-2.16 (m, 2), 2.19 (s, 3, NCH_3) 2.47 (m, 1), 2.80-2.93 (m, 2), 3.32 (m, 1, H-5), 3.51 (s, 3, OCH_3), 3.54 (m, 1, H-1), 7.01 (d, 1, $J = 7.7$, ArH), 7.28 (dd, 1, $J = 7.77$, $J = 1$, ArH), 7.60 (d, 1, $J = 1$, ArH).

Anal. Calcd for $\text{C}_{16}\text{H}_{19}\text{IN}_4\text{O}_2\cdot\text{HCl}\cdot\text{H}_2\text{O}$: C, 39.98; H, 4.61; N, 11.65. Found: C, 39.96.

Alternative synthesis for related compounds will be apparent to those of ordinary skill in the art. Additional schemes follow hereinbelow.

Synthesis

Treatment of 3β -(4-aminophenyl)tropan- 2β -carboxylic acid methyl ester (1) with the appropriate halogen gives 2. Diazotization of 2 followed by the addition of sodium azide provides the 3-halo-4-azido analog 3 (Scheme 1).


Condensation of anhydroecgonine methyl ester (4) with the appropriate acylamide oxime gives the oxadiazole 5. Addition of the appropriate aryl lithium to 5 gives the cocaine analog 6. The addition of the appropriate aryl magnesium halide to 4 gives the analog 7 (Scheme 2).

Hydrolysis of 8 gives the acid 9. Reduction of 9 with diborane gives 10. Treatment of 9 with thionyl chloride, followed by the appropriate amine gives 11. Treatment of 10

-7-


with thionyl halide or acylating agent gives 12 and 13, respectively (Scheme 3).

Scheme 1




-8-

Scheme 2



Scheme 3



-9-

Experimental2-[3-Methyl-1,2,4-oxadiazol-5-yl]-8-methyl-8-azabicyclo[3.2.1]oct-2-ene (5, R = CH₃)

Acetamide oxime (500 mg, 6.75 mmol) suspended in THF (50 mL) under nitrogen was heated at 60°C with NaH (132 mg, 5.5 mmol in oil dispersion) for 1 h. Anhydroecgonine methyl ester (2.76 mmol) and 4 Å molecular sieves (2 g) were added and the reaction mixture heated under reflux for 3 h. After cooling, the reaction mixture was filtered and the solvent removed in vacuo. The residue was chromatographed on a silica gel column eluting with CHCl₃-CH₃OH (95:5) to give the free base.

3β-Phenyl-2β-(1,2,4-oxadiazonyl-5-methyl-3-yl)-tropane (6, R = CH₃, X = H)

To an oven-dried, round-bottomed flask equipped with rubber septum and nitrogen inlet was added dry THF (25 mL) and the oxadiazole (5, R = CH₃) (261 mg, 1.27 mmol). The reaction vessel was cooled to -78°C before adding phenyl lithium (0.636 mL, 1.27 mmol) of a 2 M Et₂O solution. The reaction mixture turned dark yellow. Stirring was continued for an additional 2 h before adding brine (10 mL). The crude mixture was extracted with chloroform (3 x 45 mL), and the combined organic layers were dried (MgSO₄) and concentrated under reduced pressure to yield a yellow solid. Recrystallization from hexanes gave 53 mg (39%) of pure product as white crystals: mp 124-125°C; [α]_D +32.1° (c 0.14, MeOH); ¹H NMR (250 MHz, CDCl₃) δ 1.55-1.84 (m, 2), 1.88-1.98 (m, 2), 1.99-2.18 (m, 1), 2.24 (s, 3), 2.33 (s, 3), 2.47-2.58 (m, 1), 3.31-3.37 (t, 1, J = 7.0 Hz), 3.56-3.62 (t, 1, J = 5.7 Hz), 3.69-3.79 (ABq, 1, J = 16.1, 8.0 Hz), 4.16-4.22 (t, 1, J = 7.5 Hz), 7.05-7.14 (m, 5).

-10-

Anal. Calcd for $C_{17}H_{21}N_3O$: C, H, N.

3β -(3-Bromo-4-aminophenyl)tropane Carboxylic Acid Methyl Ester (3, X = Br)

To a round-bottomed flask containing N,N-dimethylformamide (2.5 mL) was added 3β -(4-aminophenyl)tropane carboxylic acid methyl ester (100 mg, 0.365 mmol) and N-bromosuccinimide (64.5 mg, 0.365 mmol) under a stream of nitrogen gas at ambient temperature. The resulting solution immediately turned deep red. After stirring for an additional 2.5 h, water (5 mL) was added, and the crude reaction mixture was extracted with chloroform (3 x 25 mL). The combined organic extract was dried ($MgSO_4$) and concentrated under reduced pressure to yield the product as a brown oil. Flash chromatography (5% methanol-chloroform) afforded 42 mg (33%) of pure product as a yellow oil: mp of HCl salt $194^\circ C$ dec; $[\alpha]_D -87.7^\circ$ (c 0.09, MeOH); 1H NMR (250 MHz, DMSO) δ 2.51-2.38 (m, 4), 3.39 (s, 3), 3.66-3.77 (td, 2, $J = 12.5, 2.9$ Hz), 4.17-4.59 (br s, 2), 4.67 (s, 3), 4.69-4.96 (br s, 2), 5.92 (br s, 2), 7.96-8.68 (m, 3H).

Anal. Calcd for $C_{16}H_{21}BrN_2O_2 \cdot 2 HCl \cdot 2 H_2O$: C, H, N.

General Procedure for Hydrolysis of 3β -[4-Halophenyl]tropan-2 β -carboxylic Acid Methyl Esters

The methyl ester (1.0 mmol) was dissolved in 20 mL 50% aqueous dioxane and heated to reflux. After 6 h, the solvent was evaporated and the residue crystallized from MeOH-Et₂O except as noted.

3β -[4-Iodophenyl]tropan-2 β -carboxylic Acid (9, X = I)

-11-

The starting methyl ester (0.52 mmol, 0.20 g) gave 0.137 g (71%) of the acid as a white solid: mp, 318-320°C; $[\alpha]_D^{25}$ -79.3 (c 0.55, CH₃OH); ¹H NMR (CDCl₃) δ 1.78 (m, 1), 2.02 (m, 2), 2.34 (m, 2), 2.61 (s, 3, -NCH₃), 2.7 (m, 2), 3.12 (m, 1), 3.73 (m, 2), 7.03 (d, 2, ArH), 7.62 (d, 2, ArH).

Anal. Calcd for C₁₅H₁₈INO₂: C, 48.53; H, 4.89; N, 3.77. Found: C, 48.42; H, 4.89; N, 3.71.

3β-[4-Bromophenyl]tropan-2β-carboxylic Acid (9, X = Br)

The starting ester (0.38 g, 1.1 mmol) gave 0.208 g (58%) of the acid as a white solid: mp 304-305°C; $[\alpha]_D^{25}$ -85.1° (c 0.55, CH₃OH); ¹H NMR (CDCl₃) δ 1.79 (m, 1), 2.05 (m, 2), 2.33 (m, 2), 2.65 (s, 3, -NCH₃), 2.76 (m, 2), 3.315 (m, 1), 3.77 (m, 2), 7.16 (d, 2, ArH), 7.42 (d, 2, ArH).

Anal. Calcd for C₁₅H₁₈BrNO₂: C, 55.57; H, 5.59; N, 4.32; Br, 24.65. Found: C, 55.36; H, 5.63; N, 4.28; Br, 24.53.

3β-[4-Fluorophenyl]tropan-2β-carboxylic Acid (9, X = F)

The starting methyl ester (0.60 g, 2.2 mmol) gave 0.360 g (62%) of the acid as a white solid: mp 299-300°C; $[\alpha]_D^{25}$ -92.5° (c 0.89, CH₃OH); ¹H NMR (CDCl₃) δ 1.80 (m, 1), 2.06 (m, 2), 2.36 (m, 2), 2.66 (s, 3, -NCH₃), 2.69 (m, 1), 2.79 (m, 1), 3.18 (m, 1), 3.79 (m, 2), 6.99 (m, 2, ArH), 7.25 (m, 2, ArH).

Anal. Calcd for C₁₅H₁₈FNO₂: C, 68.42; H, 6.89; N, 5.32. Found: C, 68.29; H, 6.93; N, 5.26.

3β-[4-Chlorophenyl]tropan-2β-carboxylic Acid (9, X = Cl)

The starting methyl ester (5.0 g, 6.91 mmol) gave 3.5 g

-12-

(74%) of the acid (from H₂O) as a white solid: mp 300-301°C; $[\alpha]_D^{25}$ -108.0° (c 0.10, CH₃OH); ¹H NMR (CDCl₃) δ 1.57-1.9 (m, 4), 2.25 (m, 2), 2.45 (s, 3, NCH₃), 2.52 (m, 1), 3.12 (m, 1, H-2), 3.55 (m, 2, H-1, H-5), 7.19 (dd, 4, ArH).

Anal. Calcd for C₁₅H₁₈ClNO₂ · 0.25 H₂O: C, 63.38; H, 6.56; N, 4.93. Found: C, 63.78; H, 6.56; N, 4.97.

General Procedure for Preparation of 3β-[4-Halophenyl]-2β-hydroxymethyltropane

The 2β-carboxylic acid (1.0 mmol) was suspended in dry THF (20 mL) at 0°C under N₂. A solution of BH₃ in THF (4.0 mL of 1 M solution, 4.0 mmol) was added by syringe. After 3 h, the reaction was quenched with conc. HCl (1.0 mL) and stirred for 30 min. The solvent was evaporated and the residue partitioned between dilute NH₄OH and CH₂Cl₂. The aqueous phase was further extracted with CH₂Cl₂ (3 x 50 mL). The organic extract was dried over Na₂SO₄, filtered and evaporated leaving a white solid. This was chromatographed on a silica gel flash column eluting with Et₂O-Et₃N (9:1). The sample from the column was crystallized from pentane, except as noted.

3β-[4-Iodophenyl]-2β-hydroxymethyltropane (10, X = I)

The starting 2β-carboxylic acid (0.100 g, 0.270 mmol) gave 0.055 g (57%) of the product as a white crystalline solid: mp 104-105°C; $[\alpha]_D^{25}$ -54.6 (c 0.5, CHCl₃); ¹H NMR (CDCl₃) δ 1.46 (m, 1), 1.66 (m, 1), 1.72 (d, 2), 2.17 (m, 2), 2.27 (s, 3, NCH₃), 2.48 (m, 1), 3.03 (m, 1), 3.34 (m, 2), 3.45 (m, 1), 3.75 (m, 1), 7.13 (d, 2, ArH), 7.63 (d, 2, ArH).

Anal. Calcd for C₁₅H₂₀INO: C, 50.43; H, 5.64; N, 3.92. Found: C, 50.52; H, 5.67; N, 3.84.

-13-

3 β -[4-Bromophenyl]-2 β -hydroxymethyltropane (10, X = Br)

The starting 2 β -carboxylic acid (0.150 g, 0.463 mmol) gave 0.045 g (31%) of the product as a white crystalline solid: mp 92-93°C; $[\alpha]_D^{25}$ -55.8° (c 0.5, CHCl₃); ¹H NMR (CDCl₃) δ 1.46 (m, 1), 1.62 (m, 1), 1.72 (d, 2), 2.17 (m, 2), 2.27 (s, 3, NCH₃), 2.50 (m, 1), 3.03 (m, 1), 3.34 (m, 2), 3.45 (m, 1), 3.76 (m, 1), 7.25 (d, 2, ArH), 7.43 (d, 2, ArH).

Anal. Calcd for C₁₅H₂₀BrNO: C, 58.07; H, 6.50; N, 4.52; Br, 25.76. Found: C, 57.97; H, 6.55; N, 4.45; Br, 25.83.

3 β -[4-Fluorophenyl]-2 β -hydroxymethyltropane (10, X = F)

The starting 2 β -carboxylic acid (0.263 g, 1.0 mmol) gave 0.140 g (56%) of the product as a white crystalline solid: mp 79-80°C; $[\alpha]_D^{25}$ -59.8° (c 0.5, CHCl₃); ¹H NMR (CDCl₃) δ 1.45 (m, 1), 1.63 (m, 1), 1.72 (d, 2), 2.16 (m, 2), 2.27 (s, 3, NCH₃), 2.49 (m, 1), 3.07 (m, 1), 3.34 (m, 2), 3.45 (m, 1), 3.76 (m, 1), 6.99 (m, 2, ArH), 7.32 (m, 2, ArH).

Anal. Calcd for C₁₅H₂₀FNO: C, 72.26; H, 8.08; N, 5.62. Found: C, 72.17; H, 8.10; N, 5.61.

3 β -[4-Chlorophenyl]-2 β -hydroxymethyltropane (10, X = Cl)

The starting 2 β -carboxylic acid (0.950 g, 3.4 mmol) gave 0.30 g (33%) of the product as an off-white crystalline solid: mp 104-106°C; $[\alpha]_D^{25}$ -82.4° (c 0.21, CH₃OH); ¹H NMR (CDCl₃) δ 1.45 (m, 1), 1.67 (m, 3), 2.17 (m, 2), 2.25 (s, 3, NCH₃), 2.50 (m, 1), 3.05 (m, 1, H-3), 3.30 (m, 2), 3.40 (m, 1, H-1), 3.72 (dd, 1), 7.29 (m, 4, ArH).

Anal. Calcd for C₁₅H₂₀ClNO: C, 67.78; H, 7.59; N, 5.27.

-14-

Found: C, 67.63; H, 7.69; N, 5.25.

3 β -(p-Chlorophenyl)-2 β -acetoxymethyltropane (13, X = Cl)

To a flask containing acetic anhydride (10 mL) in dry pyridine (5 mL) at ambient temperature was added 3 β -(p-chlorophenyl)-2 β -hydroxymethyltropane (95 mg, 0.32 mmol). The reaction mixture was maintained at room temperature for 2 h before diluting with water (10 mL) and adjusting the pH of the aqueous phase to 14. After extraction of the aqueous phase with chloroform (3 x 25 mL), the organic layers were combined, dried (MgSO₄) and concentrated under reduced pressure to yield the crude product as a yellow oil. Flash chromatography (CHCl₃-MeOH, 9:1) yielded 45 mg (41%) of pure product as a colorless oil: mp of HCl salt 202°C dec; [α]_D -57.1° (c 0.070, MeOH); ¹H NMR (250 MHz, CDCl₃) δ 2.02 (s, 3), 2.17-2.59 (m, 6), 2.87 (s, 3), 3.49-3.69 (m, 2), 3.99-4.22 (m, 4), 7.27-7.41 (m, 4).

Anal. Calcd for C₁₇H₂₂ClNO₂ · HCl · 0.25 H₂O: C, H, N.

3 β -(p-Chlorophenyl)-2 β -(N-methylcarbamoyl)tropane (11, R = CH₃, R₂ = H, X = Cl)

To a flask containing thionyl chloride (10 mL, _____) at 0°C was added 3 β -(p-chlorophenyl)tropane-2 β -carboxylic acid (183 mg, 0.0715 mmol). The mixture was maintained at 0°C for 4 h before concentrating under reduced pressure. The brown residue was dissolved in methylene chloride (10 mL) and cooled to 0°C before adding methylamine (5 mL). Stirring was continued for 15 min after which the excess methylamine was allowed to evaporate. The brown residue was diluted with water (25 mL) and extracted with CHCl₃ (3 x 25 mL). The combined extracts were dried (MgSO₄) and concentrated under

-15-

reduced pressure to give the crude product as a brown oil. Flash chromatography (CHCl_3 -MeOH, 9:1) yielded 72 mg (37%) of pure product as a yellow oil: mp HCl salt 138°C ; $[\alpha]_D -96.9^\circ$ (c 0.170, MeOH); ^1H NMR (250 MHz, CDCl_3) δ 1.55-1.88 (m, 5), 2.07-2.28 (m, 2), 2.31 (s, 3), 2.35-2.55 (m, 1), 2.69 (s, 3), 3.11-3.33 (m, 1), 3.40-3.49 (br s, 1), 7.14-7.26 (m, 4).

Anal. Calcd for $\text{C}_{16}\text{H}_{21}\text{ClN}_2\text{O} \cdot \text{HCl} \cdot 0.75 \text{H}_2\text{O}$: C, H, N.

3β -(p-Chlorophenyl)- 2β -chloromethyltropane (12, X = Y = Cl)

To a flask containing thionyl chloride (5 mL, _____) was added 3β -(p-chlorophenyl)- 2β -hydroxymethyltropane (64 mg, 0.24 mmol). The reaction mixture was maintained at reflux for 2 h before carefully diluting with water and adjusting the pH of the aqueous phase to 14 with conc. ammonium hydroxide. The aqueous layer was extracted with CHCl_3 (3 x 25 mL). The organic layers were combined, dried (MgSO_4), and concentrated under reduced pressure to yield the crude product as a brown oil. Flash chromatography (CHCl_3 -MeOH, 9:1) yielded 33 mg (52%) of pure product as a colorless oil: mp of HCl salt 208°C ; $[\alpha]_D -63.9^\circ$ (c 0.155, MeOH); ^1H NMR (250 MHz, CDCl_3) δ 1.05-2.50 (m, 6), 2.69 (s, 3), 2.88-3.16 (m, 2), 3.25-3.52 (m, 1), 3.78-3.89 (br s, 1), 4.02-4.15 (br s, 1), 4.55 (t, 1, J = 12.3 Hz), 7.01-7.59 (m, 4).

Anal. Calcd for $\text{C}_{15}\text{H}_{19}\text{Cl}_2\text{N} \cdot \text{HCl}$: C, H, N.

3β -(3,4-Dichlorophenyl)- 2β -chloromethyltropane (7, X = Y = Cl)

To a three-neck, round-bottomed flask containing freshly distilled ether (125 mL) and magnesium turnings (268 mg, 11.0

-16-

mmol) was added 3,4-dichloriodobenzene (2.26 g, 8.27 mmol). After 2 h, the reaction flask was equipped with a mechanical stirrer, and the Grignard reagent was cooled to -55°C before adding anhydroecgonine methyl ester (500 mg, 2.75 mmol). The resulting solution was stirred for an additional 2.5 h before being cooled to -78°C . After 1 h, 2 mL of trifluoroacetic acid was added to the solution followed by 2 h of stirring. The quenched reaction mixture was then diluted with 1 N HCl (100 mL) and extracted with ether (3 x 100 mL). The ethereal layers were discarded, and the aqueous layer was basified with conc. ammonium hydroxide and then extracted with chloroform (3 x 50 mL). The combined organic layers were dried (MgSO_4) and concentrated under reduced pressure to yield the crude product as a colorless oil. Flash chromatography (ether-triethylamine, 9:1) yielded 71 mg (9.0%) of pure product: ^1H NMR (250 MHz, CDCl_3) δ 1.52-1.76 (m, 2), 1.81-1.95 (m, 2), 1.96-2.22 (m, 2), 2.38 (s, 3), 3.07-3.15 (br s, 2), 3.21-3.32 (br s, 1), 3.45-3.65 (m, 1), 3.50 (s, 3), 7.10-7.38 (m, 3).

Anal. Calcd for $\text{C}_{16}\text{H}_{19}\text{Cl}_2\text{NO}_2 \cdot \text{HCl}$: C, H, N.

3 β -(4-Chloro-3-methylphenyl)-2 β -chloromethyltropane (7, X = Cl, Y = CH_3)

To a three-neck, round-bottomed flask containing freshly distilled ether (125 mL) and magnesium turnings (200 mg, 8.25 mmol) was added 4-chloro-3-methylbromobenzene (1.69 g, 8.25 mmol). After 2 h, the reaction flask was equipped with a mechanical stirrer, and the Grignard reagent was cooled to -55°C before adding anhydroecgonine methyl ester (500 mg, 2.75 mmol). The resulting solution was stirred for an additional 2.5 h before being cooled to -78°C . After 1 h, 2 mL of trifluoroacetic acid was added to the solution followed by 2 h of stirring. The quenched reaction mixture was then diluted

-17-

with of 1 N HCl (100 mL) and washed with ether (3 x 100 mL). The aqueous layer was basified with conc. ammonium hydroxide and extracted with CHCl_3 (3 x 50 mL). The combined organic layers were dried (MgSO_4) and concentrated under reduced pressure to yield the crude product as a colorless oil. Flash chromatography (ether-triethylamine, 9:1) yielded 45 mg (5.0%) of pure product: ^1H NMR (250 MHz, CDCl_3) δ 1.51-1.83 (m, 2), 1.97-2.21 (m, 2), 2.20 (s, 3), 2.45-2.59 (td, 1, $J = 9.5$, 2.6 Hz, 2.82-3.02 (m, 3), 3.34-3.40 (br s, 1), 3.51 (s, 3), 3.52-3.61 (br s, 1), 7.00-7.23 (m, 3).

Anal. Calcd for $\text{C}_{17}\text{H}_{22}\text{ClNO}_2 \cdot \text{HCl} \cdot 2 \text{H}_2\text{O}$: C, H, N.

3 β -(3'-Methyl-4'-fluorophenyl)tropan-2 β -carboxylic Acid Methyl Ester (7, X = F, Y = CH_3)

The title compound was prepared by modification of a reported procedure used to prepare other similar compounds.^{ref} Thus, using anhydroecgonine methyl ester (500 mg, 2.76 mmol) and 3-methyl-4-fluorophenyl magnesium bromide (prepared from 200 mg of magnesium metal and 1 mL of 3-methyl-4-fluoro-1-bromobenzene) yielded 234 mg (29%) of the title compound. The hydrochloride salt had mp 163-165°C; $[\alpha]_D^{25} -103.8^\circ$ (c 0.08, MeOH); ^1H NMR of free base of 41 (250 MHz, CDCl_3) δ 1.67 (m, 3), 2.15 (m, 2), 2.19 (s, 3, CH_3), 2.20 (s, 3, NCH_3), 2.55 (m, 2), 2.87 (m, 1, H-2), 2.93 (m, 1, H-3), 3.35 (m, 1, H-5), 3.49 (s, 3, OCH_3), 3.55 (m, 1, H-1), 6.85, 6.97 (m, 3, C_6H_3).

Anal. Calcd for $\text{C}_{17}\text{H}_{23}\text{ClFNO}_2 \cdot 1.5 \text{H}_2\text{O}$: C, 57.54; H, 7.39; N, 3.95. Found: C, 57.88; H, 7.21; N, 4.20.

[^3H]WIN 35,428 Radioligand Binding

Rat striata from male Sprague-Dawley rats (250-350 g)

-18-

were rapidly dissected, frozen, and stored at -70°C until used. The frozen rat striata were homogenized in 20 volumes of 10 mM phosphate buffer (pH 7.4) containing 0.32M sucrose using a polytron (setting 6) for 10 sec. The homogenate was centrifuged for 10 min at 50,000 x g, the resulting pellet was washed in buffer, recentrifuged, and resuspended to a tissue concentration of 10.0 mg/mL. Binding assays were carried out in a total volume of 0.5 mL containing 0.5 nM [^3H]WIN 35,428 and 1.0 mg tissue. The suspensions were incubated for 2 h on ice. Incubations were terminated by filtration with three 5 mL washes through Whatman GF/B filters previously soaked in 0.05% polyethylenimine using a Brandel M48R filtering manifold (Brandel Instruments, Gaithersburg, MD). Radioactivity was counted in 5 mL of scintillation cocktail in a Beckman LS 3801 liquid scintillation counter with an efficiency of approximately 50%. Nonspecific binding of [^3H]WIN 35,428 was defined by the presence of 30 μM (-)-Cocaine. Under these conditions, nonspecific binding was approximately 5-8% of total binding. IC_{50} values were determined from competition curves of 10-12 points utilizing the curve fitting program EBDA. Mean values and standard errors were calculated from 3-4 assays for each test drug.

Tissue Preincubation with Irreversible Agents

Tissue was prepared as described above, and the final homogenate was incubated for 60 min with either drug or vehicle as control for 60 min on ice in the above buffer. Following the 60 min incubation period, all compounds containing an azido group were then exposed to UV light (2800 Å) for 40 sec. The incubation of all compounds was terminated by centrifugation at 50,000 x g for 10 min. The resulting pellet was resuspended to a concentration of 10 mg/mL, and an aliquot was removed (0 washes). This procedure was repeated

-19-

for a total of 3 washes. Residual [^3H]WIN 35,428 binding was determined as described above. Data are expressed as the percent of specific control binding.

Testing of various compounds within the described class has given remarkably high binding values. Thus, as reported in the parent applications, receptor binding activity can be determined by degree of inhibition of the binding of [^3H]WIN 35,428. In such assays, the ligand of interest is assigned a IC_{50} value, when incubated in a 10 nM phosphate buffer, pH 7.4, containing 0.32% sucrose, with 0.5 nM [^3H]WIN 35,428 for a two hour period of time. After that, the radioactivity bound to the substrate is measured. As reported in U.S. Application Serial No. 07/564,755, on binding to a dopamine transporter receptor site, cocaine gave a IC_{50} of 89.00 nM, WIN 35,428 gave a value of 14.00 nM and a compound representative of the subject matter claimed in that application, 3 β -[4-iodophenyl]-tropane-2 β -carboxylic acid methyl ester tartrate gave a IC_{50} value of 0.25 nM. In similar assays, values of 1.35 and 4.93 nM were obtained for compounds within the class set forth above, particularly, those compounds bearing a carboxylic acid moiety, or a heterocyclic moiety. Similar values may be obtained for the remaining members of the class.

When bearing an appropriate radioactive label, such as ^{11}C or ^{123}I , ^{125}I or ^{131}I , these compounds, preferential binders to dopamine transporter and serotonin transporter binding sites, can be used as imaging agents for both positron emission tomography (PET) as well as single photon emission computed tomography (SPECT). PET may require the [^{11}C] labeled form of the drug, while radioactive iodine-labeled compounds may be used in SPECT scanning.

As noted, such scanning has a variety of utilities. The

-20-

actual density and distribution of cocaine receptors in various parts of the brain and CNS is of interest, and can be mapped through the use of these compounds. Additionally, as noted above, identification of degeneration of nerve terminals, corresponding to a loss of dopamine transporter sites, can be determined by this scanning, to diagnose Parkinson's Disease. In addition, progression of the disease, and efficacy of treatment, can be monitored by such scanning. Similarly, degeneration of specific nerves in the brain due to exposure to various toxins can be detected or monitored by the scanning made possible by these compounds.

As an additional use, drugs having high affinity for the transporter sites bound to by these compounds, particularly serotonin and dopamine transporter sites, can be screened, using these compounds, in the identified scanning methods. The scanning itself is conventional, given the use of these compounds, and does not constitute an aspect of the invention per se. Affinity values for representative compounds are given in the following table.

-21-

Table. Potencies of Cocaine and Analogs in Inhibiting Binding of [³]-3 β -(4-Fluorophenyl)tropan-2 β -carboxylic Acid Methyl Ester (WIN 35,428)

Compound	IC ₅₀ (nM)
Cocaine	102
2 (X = I)	1.35
3 (X = I)	4.93
6 (R = CH ₃ , X = H)	48
7 (X = Y = Cl)	0.79
7 (X = Cl, Y = CH ₃)	0.81
9 (X = Br)	279
9 (X = I)	474
9 (X = Cl)	2070
9 (X = F)	2740
10 (X = Br)	1.49
10 (X = Cl)	1.53
10 (X = I)	2.2
10 (X = F)	47.3
11 (R ₁ = CH ₃ , R = H, X = Cl)	12.4
12 (X = Y = Cl)	2.64
13 (R = CH ₃ , X = Cl)	1.6

This invention has been described in both generic terms, and by reference to specific description. No specific


-22-

description or example is considered binding, unless so identified. Alternate forms and methods will occur to those of ordinary skill in the art, without the exercise of inventive faculty, and remain within the scope of this invention, save as limited by the claims set forth below.


-23-

Claims

1. A compound of the formula below,



wherein Y = CH₂R₃, CO₂R₂ or



R₁ = hydrogen, C₁₋₅ alkyl,

R₂ = hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₄ alkoxy, C₁₋₆ alkynyl, halogen or amine,

R₃ = OH, hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₄ alkoxy, Cl, Br, I, CN, NH₂, NHC₁₋₆ alkyl, NC₁₋₆ alkyl, OCOC₁₋₆ alkyl, OCOC₁₋₃ alkylaryl,

A = S, O or NH

X = H, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₄ alkoxy, C₁₋₆ alkynyl, halogen, amino, acylamido,

and

Z = H, I, Br, Cl, F, CN, CF₃, NO₂, N₃, OR₁, CO₂NH₂, CO₂R₁, C₁₋₆

-24-

alkyl, NR_4R_5 , NHCOR_5 , NHCO_2R_6 ,

wherein $\text{R}_4\text{-R}_6$ are each C_{1-6} alkyl and wherein said compound bears at least one iodine or carbon atom which is radioactive.

2. The compound of Claim 1, wherein said radioactive atom is [^{11}C].

3. The compound of Claim 1, wherein said compound bears an iodine atom, and said iodine atom is radioactive, and selected from the group consisting of ^{123}I , ^{125}I and ^{131}I .

INTERNATIONAL SEARCH REPORT

PCT/US92/09482

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :A61K 43/00; A61K 49/02; C07D 451/00; C07D 451/02

US CL :424/1.1,546/124,546/125,546/132

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. :

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US,A, 4,179,567 (CLARKE ET AL) See col. 1, line 40, to col. 2, line 2	1-3
X Y	US,A, 5,128,118 (CARROLL ET AL) See abstract and col. 1, line 65 to col 2, line 46	<u>1 and 3</u> 2
X	US,A, 3,813,404 (CLARKE ET AL) See col. 1, line 10 to line 68	1-2

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

04 JANUARY 1993

Date of mailing of the international search report

19 JAN 1993

Name and mailing address of the ISA/
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231Authorized officer *Matthew Zmurko*
MATTHEW ZMURKO

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0511

Form PCT/ISA/210 (second sheet)(July 1992)*

